

A Proposal of Residual Reduction method with preconditioning based on IDR Theorem

List of authors:

Seiji Fujino¹

Takashi Sekimoto² and Yusuke Onoue²

IDR(s) method [4] and extended variants of IDR(s) method [3] based on the IDR Theorem have been proposed one after another. The authors also proposed IDR(s)-based iterative method with iteration matrix of SOR (Successive Over-Relaxation) method in order to accelerate its convergence [2]. In my talk, we propose an algorithm of IDR(s)-based Residual Reduction method using precondition with Eisenstat trick [1]. We refer to IDR(s)-R2 method. We present summary of performance of IDR(s)-R2 method using some preconditions with Eisenstat trick in Table 1. We realize that preconditioned IDR(s)-R2 method outperforms with the conventional iterative methods.

Table 1: Summary of performance of preconditioned IDR(s)-R2 method.

method	precond.	no. of convergence	no. of nonconvergence	no. of fastest case	total (%)
IDR(s)-R2	none	11/18	7/18	0/18	0
	Gauss-Seidel-L	18	0	8	14 (61%)
	Eisenstat-L	17	1	1	
	Eisenstat-UL	17	1	5	
BiCGStab	ILU(0)	15	3	5	5(22%)
BiCGSafe	ILU(0)	15	3	4	4(17%)

References

- [1] Eisenstat, S. C.: Efficient implementation of a class of preconditioned conjugate gradient methods, SIAM J. Sci. Stat. Compute., **2**(1981), 1-4.
- [2] Fujino, S., Sonneveld, P. and van Gijzen, M.B. and Onoue, Y., Application of the IDR Theorem to Stationary Iterative Methods and their Performance Evaluation, The Abstract of SIAM LA09, Monterey (2009).
- [3] Kusakabe, Y., Fujino, S., A proposal of Jacobi method based on extended Induced Dimension Reduction Theorem aimed at high convergence rate, The Proc. of EMAC2009, Adelaide (2009).
- [4] Sonneveld, P., van Gijzen, M.B., IDR(s): a family of simple and fast algorithms for solving large nonsymmetric linear systems, SIAM J. Sci. Comput., **31**(2008), 1035–1062.

¹Research Institute for Information Technology, Kyushu University

²Graduate School of Information Science and Electrical Engineering, Kyushu University